

## ARTICLE

# Influence of exogenous salicylic acid on antioxidant enzyme activities in the roots of salt stressed tomato plants

Ágnes Szepesi\*, Péter Poór, Katalin Gémes, Edit Horváth, Irma Tari

Department of Plant Biology, University of Szeged, Szeged, Hungary

**ABSTRACT** Up-regulation of the antioxidant system provides protection against NaCl-induced oxidative damage in plants. Activities of antioxidant enzymes and hydrogen peroxide ( $H_2O_2$ ) levels in the roots of tomato (*Lycopersicon esculentum* Mill. L.) plants were investigated to assess the antioxidant protection offered by long-term (3 weeks) exogenous salicylic acid (SA) pre-treatment against 100 mM NaCl induced salt stress. Our results suggest that  $10^{-4}$  M SA was effective in increasing the activities of enzymes involved in the antioxidant system.

**Acta Biol Szeged 52(1):199-200 (2008)****KEY WORDS***Lycopersicon esculentum* Mill. L.  
oxidative stress  
salicylic acid  
salt stress

When plants are subjected to saline conditions, reactive oxygen species (ROS) like superoxide, hydrogen peroxide ( $H_2O_2$ ) and hydroxyl radicals are generated (Dat et al. 2000; Farooq and Azam 2007). Salinity imposes oxidative stress on tissues that seriously disrupt normal metabolism of plants through oxidation of membrane lipids, proteins and nucleic acids in the absence of protective mechanisms (Noctor and Foyer 1998; Hernández et al. 2001). In order to avoid the harmful effects of ROS, plants evolved an effective scavenging system composed of enzymatic antioxidants, such as superoxide dismutase (SOD, EC 1.15.1.1), peroxidase (POD, EC 1.11.1.7) and catalase (CAT, EC 1.11.1.6), a  $H_2O_2$  generating and degrading enzymes, respectively (Hajiboland and Hasami 2007).

Salicylic acid (SA) is considered as a hormone-like substance that has important role in the regulation of plant growth and development. There is strong evidence that SA mediates the oxidative burst. SA specifically binds to catalase and inhibits the activity of the enzyme (Chen et al. 1993). Thus, exogenous SA can increase  $H_2O_2$  content of tissues, can induce the expression of antioxidant enzymes and increase plant tolerance to the abiotic stressors. In the present work salt stress acclimation of tomato plants, which had previously been grown for three weeks in low concentrations of SA ( $10^{-7}$  M and  $10^{-4}$  M), was investigated in order to follow the changes in  $H_2O_2$  contents of tissues.

## Materials and Methods

Tomato (*Lycopersicon esculentum* Mill. cvar Rio Fuego) plants were grown hydroponically in a greenhouse under  $180 \mu\text{mol m}^{-2} \text{s}^{-1}$  light intensity and at 12/12 day/night photoperiod.

The temperature was maintained at 25°C and the relative humidity was 55-60%. The plants were pre-treated with  $10^{-7}$ - $10^{-4}$  M SA for three weeks. The salt stress was induced by 100 mM NaCl treatment for 1 week. Enzyme activities were measured after 1 week of salt stress. 750 mg of plant tissue was homogenized in 3 ml extraction buffer (50 mM phosphate buffer pH 7.0, containing 1 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF) and 1% polyvinyl-polypyrrolidone). After centrifugation the supernatant was used for enzyme activity assays. Superoxide dismutase (SOD) activity was determined by measuring the ability of the enzyme to inhibit the photochemical reduction of nitroblue tetrazolium (NBT) in the presence of riboflavin in light (Dhindsa et al. 1981). Catalase (CAT) activity was determined by the decomposition of  $H_2O_2$  and was measured spectrophotometrically by following the decrease in absorbance at 240 nm (Upadhyaya et al. 1985). Guaiacol peroxidase (POD) activity was determined by monitoring the increase in absorbance at 470 nm during the oxidation of guaiacol (Upadhyaya et al. 1985). The protein contents of the extracts were determined by the method of Bradford (1976). Histochemical identification of hydrogen peroxide by 3,3'-diaminobenzidine (DAB) staining occurred according to Thordal-Christensen et al. (1997).

## Results

The first line of defense against the effects of high salinity has to be induced in the root system. In order to maintain the integrity of membranes in root cells, effective enzymatic mechanisms for scavenging ROS, among them  $H_2O_2$ , were developed in plants. The activity of a  $H_2O_2$  generating enzyme, SOD, increased after SA pre-treatments, and the activity remained much higher under salt stress than in the control plants exposed to 100 mM NaCl.

\*Corresponding author. E-mail: [szepesia@bio.u-szeged.hu](mailto:szepesia@bio.u-szeged.hu)

In parallel, increased CAT activities could be detected in the roots pre-treated with  $10^{-4}$  M SA after salt stress compared to the control. This ensured an effective decomposition of  $H_2O_2$  in root tissues. The activity of peroxidases, which may generate or scavenge  $H_2O_2$ , decreased in the roots after salt stress.

With DAB uptake by roots, hydrogen peroxide can be localized in vivo and in situ in plant tissues (Thordal-Christensen et al. 1997). Brown marks were detectable in the root tips in control plants after 100 mM NaCl treatment. However, very few brown marks emerged within the  $10^{-4}$  M SA pre-treated roots under salt stress, accordingly, long-term pre-treatment of tomato with both SA concentrations led to the degradation of  $H_2O_2$  in some degree in the root tissues.

## Discussion

ROS generated in cells are highly reactive in nature and destroy the normal cellular function and metabolism. Determining the role of various antioxidant enzymes in the salt tolerance of tomato, Mittova et al. (2002) found that higher salt tolerance of wild tomato (*Lycopersicon pennellii*) as compared to cultivated tomato (*Lycopersicon esculentum* Mill. L.) was correlated with increased activities of SOD and POD.

In agreement with these observations, our results also showed that oxidative stress occurred in the control plants exposed to high salinity. Our antioxidant enzyme activity measurements suggest that  $10^{-4}$  M SA pre-treatment induced the enzymatic antioxidant system effectively by enhancing SOD and CAT enzyme activities. High activities of SOD and CAT in wild tomato (*Lycopersicon pennellii*) correlated with higher salt tolerance and better protection against oxidative stress as compared with the salt sensitive cultivated genotype (Shalata et al. 2001).

Our data suggest that  $10^{-7}$  M SA pre-treated plants had lower capacity for scavenging ROS, generated by salt stress than  $10^{-4}$  M SA pre-treated tomatoes. The better protection by  $10^{-4}$  M SA results from the greater efficiency of the  $H_2O_2$  degradation under salt stress in the leaves and roots of pre-treated plants. These mechanisms such as reduced oxidative damage have been proposed to explain the contribution of salicylic acid to the salt tolerance of tomato plants.

## Acknowledgements

This work was supported by grants from the Hungarian National Scientific Research Foundation (OTKA TO38392) and GVOP-3.2.1.-2004-04-0419/3.0. The authors are grateful to Mrs. Kispálné Szabó Ibolya for valuable technical assistance.

## References

- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248-254.
- Chen Z, Silva H, Klessig DF (1993) Active oxygen species in the induction of plant systemic acquired resistance by salicylic acid. *Science* 262:1883-1886.
- Dat J, Vandenabeele S, Vranová E, Van Montagu M, Inzé D, Van Breusegem F (2000) Dual action of the active oxygen species during plants stress responses. *Cell Mol Life Sci* 57:779-795.
- Desikan R, A-H-Mackerness S, Hancock JT, Neill SJ (2001) Regulation of the *Arabidopsis* transcriptome by oxidative stress. *Plant Physiol* 127:159-172.
- Dhindsa RS, Plumb-Dhindsa P, Thorpe TA (1981) Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J Exp Botany* 32: 93-101.
- Farooq S, Azam F (2007) Differences in behaviour of salt tolerant and salt and water deficiency tolerant wheat genotypes when subjected to various salinity levels. *Cer Res Com* 35:63-70.
- Hajiboland R, Hasani B (2007) Responses of antioxidant defence capacity and photosynthesis of bean (*Phaseolus vulgaris* L.) plants to copper and manganese toxicity under different light intensities. *Acta Biol Szeged* 51:93-106.
- Hernández JA, Ferrer MA, Jiménez A, Barceló AR, Sevilla F (2001) Antioxidant systems and  $O_2^-/H_2O_2$  production in the apoplast of pea leaves. Its relation with salt-induced necrotic lesions in minor veins. *Plant Physiol* 127:817-831.
- Horváth E, Szalai G, Janda T (2007) Induction of abiotic stress tolerance by salicylic acid signaling. *J Plant Growth Regul* 26:290-300.
- Mittova V, Tal M, Volokita M, Guy M (2002) Salt stress induces upregulation of an efficient chloroplast antioxidant system in the salt-tolerant wild tomato species *Lycopersicon pennellii* but not in the cultivated species. *Physiol Plant* 115:393-400.
- Noctor G, Foyer CH (1998) Ascorbate and glutathione: keeping active oxygen under control. *Annu Rev Plant Physiol Plant Mol Biol* 49:249-279.
- Shalata A, Mittova V, Volokita M, Guy M, Tal M (2001) Response of the cultivated tomato and its wild salt-tolerant relative *Lycopersicon pennellii* to salt-dependent oxidative stress: The root antioxidative system. *Physiol Plant* 112:487-494.
- Thordal-Christensen H, Zhang Z, Wei Y, Collinge DB (1997) Subcellular localization of  $H_2O_2$ . *The Plant Journal* 11:1187-1194.
- Upadhyaya A, Sankhla D, Davis TD, Sankhla N, Smith BN (1985) Effect of paclobutrazol on the activities of some enzymes of activated oxygen metabolism and lipid peroxidation in senescing soybean leaves. *J Plant Physiol* 121:453-461.